

<p><b>DTaP</b></p> <p><b>DT:</b> for children who can't have pertussis vaccine</p>	<p><b>Diphtheria</b> <b>Tetanus</b> (acellular) <b>Pertussis</b></p>	<p>Received at <b>6 weeks through 6 years</b></p>	<p><b>4-5 doses</b> Children need 5 doses. Some will only get 4 doses because they got behind in getting their vaccines</p>	<p>Brand name of vaccine include: <b>DTaP:</b> Infanrix, Daptacel, Tripedia, Acel-Imune</p> <p><b>DTaP+other vaccine</b></p> <ul style="list-style-type: none"> <li>• Pentacel (DTaP+IPV+Hib)</li> <li>• Pediarix (DTaP+IPV+Hep B)</li> <li>• Kinrix (DTaP+IPV)</li> <li>• TriHIBit (DTaP+Hib)</li> </ul>
<p><b>Td</b></p>	<p><b>Tetanus</b> <b>diphtheria</b></p>	<p>Received at <b>7 years of age and older</b></p>	<p><b>1-3 doses</b> Most children will not get any doses because they got enough DTaP vaccine</p>	<p>Brand name of <b>Td</b> vaccine include:</p> <ul style="list-style-type: none"> <li>• Td (generic)</li> <li>• Decavac</li> <li>• Tenivac</li> </ul>
<p><b>Tdap</b></p>	<p><b>Tetanus</b> <b>diphtheria</b> (acellular) <b>pertussis</b></p>	<p>Received at <b>11 years of age and older</b> Using specific rules, MCIR will count some doses given at 10 years of age or younger</p>	<p><b>1 dose</b> Persons should only get one dose of this vaccine</p>	<p>Brand name of <b>Tdap</b> vaccine include:</p> <ul style="list-style-type: none"> <li>• Adacel</li> <li>• Boostrix</li> </ul>
<ul style="list-style-type: none"> <li>• Upper case "<b>T</b>" means there is about the same amount of tetanus in DTaP, Tdap and Td.</li> <li>• Upper case "<b>D</b>" and "<b>P</b>" means there is more diphtheria and pertussis in DTaP than in Tdap and Td; lower case letters ("d" "p") means there is less.</li> <li>• <b>DTaP</b> produces fewer side effects and is a safer version of an older vaccine called <b>DTP</b>, which is no longer used in the United States. <b>DTP:</b> P=whole cell pertussis</li> </ul>				